

Developing resources for improved chronic kidney disease management in primary care

Brook Butler


Knowledge Transfer Associate

Greater Manchester CLAHRC



IMPACT

Chronic Kidney Disease (CKD)



■ Death ■ Cardiovascular event ■ Hospitalisation



- Affects about 6% of adults in the UK
- May progress to end stage renal failure, requiring transplant or dialysis
- Greatly increases a person's risk of suffering a stroke, heart attack or death
- **BUT** often undiagnosed and poorly managed



The GM CLAHRC CKD Collaborative

The GM CLAHRC CKD Collaborative was a 12 month improvement project, which had two aims:

- 1. Halve the prevalence gap**
- 2. Ensure that 75% of all patients are treated to the NICE recommended blood pressure targets**




LNR CLAHC's PSP-CKD Study

Aims:

- To determine whether reinforcement of best practice in the management of key aspects of CKD care improves clinical outcomes
- To improve coding of CKD and prevalence on chronic disease registers
- To increase interest in and capacity for primary care research in Northamptonshire
- To implement and evaluate a new model of partnership working between primary and secondary care



The Challenge in LNR: Average Prevalences 08-11



Population of CKD registers has plateaued



The LNR Audit Tool

A MiQuest based tool, which addresses:

- Uncoded patients
- Accuracy of existing coding
- Risk stratification
- Practice audit
- Referral management



Results:

Uncoded patients

1	Patients not coded as CKD				
2	Identifier	CKD confirmed by existing eGFR data?	Suggested Classification	Evidence: Latest eGFR<60	Evidence: Proteinuria
187		eGFR data confirms CKD3 at least	CKD Stage 3A	57 on 07/02/2012	ACR <30mg/mmol
188		eGFR data confirms CKD3 at least	CKD Stage 3A	48 on 01/02/2012	without proteinuria on dipstick
189		eGFR data confirms CKD3 at least	CKD Stage 3A	51 on 09/01/2012	without proteinuria on dipstick
190		eGFR data confirms CKD3 at least	CKD Stage 3A	49 on 13/01/2012	ACR <30mg/mmol
191		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	49 on 22/01/2007	without proteinuria on dipstick
192		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	52 on 22/01/2007	without proteinuria on dipstick
193		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	58 on 22/01/2008	without proteinuria on dipstick
194		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	55 on 10/11/2011	without proteinuria on dipstick
195		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	58 on 20/11/2007	ACR
196		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	56 on 30/08/2007	without proteinuria on dipstick
197		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	56 on 09/11/2009	without proteinuria on dipstick
198		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	53 on 08/01/2008	without proteinuria on dipstick
199		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	59 on 08/04/2011	ACR <30mg/mmol
200		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	59 on 06/04/2011	without proteinuria on dipstick
201		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	58 on 16/09/2011	without proteinuria on dipstick
202		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	57 on 14/08/2007	without proteinuria on dipstick
203		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	59 on 03/03/2011	without proteinuria on dipstick
204		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	53 on 14/03/2007	without proteinuria on dipstick
205		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	58 on 02/03/2009	PCR <50mg/mmol
206		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	57 on 22/09/2011	without proteinuria on dipstick
207		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	59 on 08/09/2011	without proteinuria on dipstick
208		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	52 on 07/06/2011	ACR <30mg/mmol
209		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	57 on 13/03/2007	without proteinuria on dipstick
210		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	58 on 30/11/2010	ACR <30mg/mmol
211		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	57 on 02/03/2011	without proteinuria on dipstick
212		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	55 on 01/11/2010	without proteinuria on dipstick
213		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	53 on 10/01/2012	No proteinuria data
214		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	54 on 17/02/2011	without proteinuria on dipstick
215		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	57 on 06/09/2011	without proteinuria on dipstick
216		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	58 on 26/02/2008	without proteinuria on dipstick
217		eGFR data confirms CKD3 at least	CKD Stage 3A but later eGFRs exist over 6	57 on 25/06/2007	without proteinuria on dipstick

Results:

Accuracy of existing coding

1	Accuracy of existing CKD coding				
2	Identifier	Coded Classification	Classification by latest data	Evidence: Latest eGFR<60	Evidence: Proteinuria
362		CKD stage 5	CKD Stage 4	16 on 03/01/2012	without proteinuria on dipstick
363		CKD stage 3B	CKD Stage 4	23 on 11/11/2011	without proteinuria on dipstick
364		CKD stage 4	CKD Stage 4	18 on 27/01/2012	without proteinuria on dipstick
365		CKD stage 4	CKD Stage 4	29 on 20/09/2011	ACR <30mg/mmol
366		CKD stage 5	CKD Stage 4	23 on 15/12/2011	ACR <30mg/mmol
367		CKD stage 4	CKD Stage 4	23 on 22/09/2011	ACR <30mg/mmol
368		CKD stage 4	CKD Stage 4	29 on 09/06/2009	PCR <50mg/mmol
369		CKD stage 4	CKD Stage 4 equivocal proteinuria	24 on 17/01/2012	+ protein on dipstick
370		CKD stage 4	CKD Stage 4 with proteinuria	16 on 15/07/2011	PCR >=50mg/mmol
371		CKD stage 4 with proteinu	CKD Stage 4 with proteinuria	27 on 09/05/2011	PCR >=50mg/mmol
372		CKD stage 4	CKD Stage 4 with proteinuria	29 on 20/09/2011	ACR >=30mg/mmol
373		CKD stage 4	CKD Stage 4 with proteinuria	28 on 25/11/2011	+++ protein on dipstick
374		CKD stage 3	CKD Stage 5	5 on 12/07/2011	without proteinuria on dipstick
375		CKD stage 4	CKD Stage 5	14 on 06/06/2011	ACR <30mg/mmol
376		CKD stage 4	CKD Stage 5 with proteinuria	12 on 17/01/2012	ACR >=30mg/mmol
377		CKD stage 4	CKD Stage 5 with proteinuria	13 on 03/11/2011	ACR >=30mg/mmol
378		CKD stage 3	Not enough eGFR data to confirm sta	59 latest<60 on 11/11/2011	
379		CKD stage 3	Not enough eGFR data to confirm sta	56 latest<60 on 17/01/2007	
380		CKD stage 3A	Not enough eGFR data to confirm sta	59 latest<60 on 23/01/2012	
381		CKD stage 2	Not enough eGFR data to confirm sta	57 latest<60 on 31/07/2006	
382		CKD stage 3A	Not enough eGFR data to confirm sta	51 latest<60 on 22/12/2011	
383		CKD stage 3	Not enough eGFR data to confirm sta	33 latest<60 on 19/05/2009	
384		CKD stage 3	Not enough eGFR data to confirm sta	36 latest<60 on 29/01/2010	
385		CKD stage 3	Not enough eGFR data to confirm sta	58 latest<60 on 16/01/2007	
386		CKD stage 3	Not enough eGFR data to confirm sta	42 latest<60 on 25/05/2011	
387		CKD stage 3	Not enough eGFR data to confirm sta	56 latest<60 on 10/07/2006	
388		CKD stage 3	Not enough eGFR data to confirm sta	53 latest<60 on 17/07/2006	
389		CKD stage 3	Not enough eGFR data to confirm sta	53 latest<60 on 15/11/2006	
390		CKD stage 2	Not enough eGFR data to confirm sta	54 latest<60 on 03/09/2007	
391		CKD stage 3	Not enough eGFR data to confirm sta	59 latest<60 on 12/01/2007	

Results:

Risk register

1											
2											
3	175	16	238	111	11	531					82
4	Diabetic	Urinary outflow tract obstruct	Cardiovasc Disease	Smoker	Black / Asian Ethnicity	Ranked by combined risk score	On NSAID				
5	Identifier	Identifier	Identifier	Identifier	Identifier	Identifier	Score	Identifier	Code		
6								7	j22a.		
7								7	j282.		
8								6	j282.		
9								6	j22y.		
10								6	j2bx.		
11								6	j282.		
12								6	j282.		
13								5	j22e.		
14								5	j282.		
15								5	j2c2.		
16								5	j2c2.		
17								5	j22o.		
18								5	j22o.		
19								5	j2n3.		
20								5	j28Y.		
21								5	j2c1.		
22								5	j2c2.		
23								5	j2c1.		
24								5	j22e.		
25								5	j282.		
26								5	j2c2.		
27								4	j282.		
28								4	j2n3.		
29								4	j282.		
30								4	j281.		

7	j22a.
7	j282.
6	j282.
6	j22y.
6	j2bx.
6	j282.
6	j282.
5	j22e.
5	j282.
5	j2c2.

High Risk Patients

Results:

Practice audit

CKD Audit Data							
Practice Code	C12345	All these patients have either eGFR <60 or other code indicating renal impairment					
Date of data extraction	20120217						
Total population over 17 at this date	8968						
Registers and process measures							
	CKD1	CKD2	CKD3	CKD4	CKD5	Uncoded	RRT
Number with at least 1 eGFR<60 (uncoded) or coded CKD (excluding RRT)		30	335	21	3	248	11
Observed % prevalence for population over 17	0.0	0.3	3.7	0.2	0.0	2.8	0.1
Exp % prev >17 adj by age /sex (CKD3-5 only)			5.7	0.3	0.1		
Patients with coded CKD diagnosis->							
	Diabetes	No Diabetes	Proteinuria	No Proteinuria	Unassessed		
Number with coded CKD (any stage)	117	272	46	257	86		
% of population over 17 with coded CKD	1.3	3.0	0.5	2.9	1.0		
Blood Pressure							
	Diabetes	No Diabetes	Proteinuria	No Proteinuria	Overall		
Number with BP recorded in last year	115	253	43	249			
% with BP recorded in last year	98.3	93.0	93.5	96.9	75.1		
Number treated to target using latest BP in last year	35	165	13	162			
% BP treated to target	29.9	60.7	28.3	63.0	45.0		
Proteinuria and ACE/ARB usage							
Number of DM with ACR recorded ever; No DM ACR/PCR recorded ever	114	200					
% with proteinuria assessed ever	97.4	73.5					
Number of DM ACR>2.5 m, >3.5 f; No DM (ACR>70/PCR>100 OR ACR>30/PCR>50 & HBP)	36	9					
% prev microalbuminuria in diabetes / ACE/ARB Rxable proteinuria in non-diabetes	30.8	3.3					
Number on ACE/ARB in above groups	24	6					
% on ACE/ARB in above groups	66.7	66.7					
Last eGFR done within recommended time frame							
	CKD1	CKD2	CKD3	CKD4	CKD5	Uncoded	
CKD1,2 in last 12m, Uncoded / CKD3A/B in last 6m, CKD4 last 3m, CKD5 in last 6w		6	131	7		33	
% of each CKD group with eGFR done within recommended time frame	#DIV/0!	20.0	39.1	33.3	0.0	13.3	
NSAID usage in CKD (NOT recommended!)							
	CKD1	CKD2	CKD3	CKD4	CKD5	Uncoded	
Number on NSAID		4	38	1		39	
% of each CKD group on an NSAID	#DIV/0!	13.3	11.3	4.8	0.0	15.7	

Results:

Referral management



1	Identifier	Alert	Intervention	Referral
2		eGFR < 45ml/min eGFR declining - expected <15 by 80yo	Check eGFR	Consider referral as eGFR decline >5 ml/min/1.73 m2 pa
3		eGFR declining - expected <15 by 80yo	Check PCR Check eGFR Review NSAID use	Consider referral as eGFR decline >5 ml/min/1.73 m2 pa
4		Cardiovascular disease	Check BP Check eGFR	Consider referral as eGFR decline >5 ml/min/1.73 m2 pa
5		eGFR < 45ml/min eGFR declining - expected <15 by 80yo	Check eGFR Review NSAID use	Consider referral as eGFR decline >5 ml/min/1.73 m2 pa
6		eGFR declining - expected <15 by 80yo Cardiovascular d	Check PCR Check eGFR	Consider referral as eGFR decline >5 ml/min/1.73 m2 pa
7		Diabetes eGFR declining - expected <15 by 80yo Cardia	Check ACR Check eGFR	Consider referral as eGFR decline >5 ml/min/1.73 m2 pa
8		High Chol>6mmol/l Diabetes eGFR declining - expected	BP-needs better control<130/80	Consider referral as eGFR decline >5 ml/min/1.73 m2 pa Consider
9		eGFR < 45ml/min ...and Hb<10.5g/dl Proteinuria presen	Check BP Consider ACE/ARB	Consider referral as high ACR
10		eGFR < 45ml/min Proteinuria present Diabetes eGFR c	BP-needs better control<130/80	Consider referral as high ACR, unless known due to their diabetes e
11		High Chol>6mmol/l Current smoker	Consider ACE/ARB Check eGFR	Consider referral as high PCR
12		Diabetes	BP-needs better control<130/80 Check eGFR	Consider referral as high PCR, unless known due to their diabetes e
13		Diabetes eGFR declining - expected <15 by 80yo Cardia	BP-needs better control<130/80	Consider referral for better BP control
14		High SBP>150mmHg High DBP>90mmHg Cardiovascul	BP-needs better control<140/90	Consider referral for better BP control
15		High SBP>150mmHg	BP-needs better control<140/90 Check eGFR Review NSAID u	Consider referral for better BP control
16		High SBP>150mmHg Diabetes Cardiovascular disease	BP-needs better control<130/80	Consider referral for better BP control
17		eGFR < 45ml/min High DBP>90mmHg Diabetes	BP-needs better control<130/80	Consider referral for better BP control
18		eGFR < 45ml/min Diabetes Urinary outflow obstruction	BP-needs better control<130/80 Review NSAID use	Consider referral for better BP control
19			BP-needs better control<140/90 Check eGFR Review NSAID u	Consider referral for better BP control
20		High SBP>150mmHg Cardiovascular disease	BP-needs better control<140/90 Check eGFR	Consider referral for better BP control
21		High SBP>150mmHg	BP-needs better control<140/90 Check eGFR	Consider referral for better BP control
22		Diabetes Cardiovascular disease Current smoker	BP-needs better control<130/80	Consider referral for better BP control
23		eGFR < 45ml/min High SBP>150mmHg Diabetes Card	BP-needs better control<130/80	Consider referral for better BP control
24		Diabetes Cardiovascular disease	BP-needs better control<130/80	Consider referral for better BP control
25		Diabetes	BP-needs better control<130/80 Check ACR	Consider referral for better BP control
26		eGFR < 45ml/min High SBP>150mmHg Diabetes eGFR	BP-needs better control<130/80 Consider ACE/ARB Check eG	Consider referral for CKD 4 or 5 patients
27		eGFR < 45ml/min ...and Hb<10.5g/dl Diabetes eGFR declining - expected <15 by 80yo Cardiovascular disease Curren		Consider referral for CKD 4 or 5 patients
28		eGFR < 45ml/min Proteinuria present Diabetes Cardio	Consider ACE/ARB Check eGFR	Consider referral for CKD 4 or 5 patients
29		eGFR < 45ml/min		Consider referral for CKD 4 or 5 patients
30		eGFR < 45ml/min High SBP>150mmHg Cardiovascular	BP-needs better control<140/90 Check eGFR	Consider referral for CKD 4 or 5 patients
31		eGFR < 45ml/min Cardiovascular disease		Consider referral for CKD 4 or 5 patients
32		eGFR < 45ml/min High Chol>6mmol/l eGFR declining - e	Check eGFR	Consider referral for CKD 4 or 5 patients
33		eGFR < 45ml/min Diabetes Cardiovascular disease	BP-needs better control<130/80 Check eGFR Review NSAID u	Consider referral for CKD 4 or 5 patients

GM CKD Phase 2: Utilising new resources



The second GM CKD Improvement Project had two new resources to aid effectiveness and efficiency:

- LNR CLAHRC's audit tool
- GM CLAHRC's Improvement Guide

A big difference in achievement was seen.



IMPAKT



IMproving Patient care and Awareness of Kidney disease progression Together



Draft screenshot of IMPAKT tool

Options ▾
Chris Abram ▾

Dashboard

Report Builder

Comparative Data


Practice Data

Upload


Support

Community


DASHBOARD




Report Builder
[Learn more](#)




Comparative User data
[Learn more](#)




Your Practice data
[Learn more](#)



Upload anonymized data and compare your results with national collective statistics
[Upload comparative data](#)



Access user guides, white papers and FAQ's through our Knowledge base
[Get help and support](#)



A forum and community for UK Healthcare Professionals with an interest in CKD
[Visit the forum](#)

NATIONAL CKD AUDIT DATA

Registers and Process Measures

	CKD1	CKD2	CKD3	CKD4	CKD5	Uncoded	RRT
Number with at least 1 eGFR<80 or coded CKD (excluding RRT)	-	3	238	18	2	383	12
% of population over 18	0.0	0.0	2.5	0.2	0.0	4.1	0.1

Last eGFR done with recommended time frame

	CKD1	CKD2	CKD3	CKD4	CKD5	Uncoded
CKD1, 2 in last 12m, Uncoded / CKD3-5 in last 6m, CKD4 last 3m, CKD5 in last 6w	-	3	114	11	1	51
% of each CKD group with eGFR done within recommended time frame	#DIV/0!	100.0	47.9	61.1	50	13.3

Blood Pressure

	Diabetes	No Diabetes
Number with BP recorded in last year	71	177
% with BP recorded in last year	99.6	99.7

Proteinuria and ACE/ARB usage

	Diabetes	No Diabetes
Number of DM with ACR recorded ever; No DM ACR/PCR recorded ever	88	101

Your CKD Audit Overview

Date of Last Data Extraction ⓘ
01-04-12

Total population over 18 at this date ⓘ
9,406

Patients with coded CKD diagnosis ⓘ
161

No. of participating practices ⓘ
64

Date of Data ⓘ
4.87

Practice Code ⓘ
XYZ

Download Practice Data



Further Information

www.impakt.org.uk

Greater Manchester CLAHRC - <http://clahrc-gm.nihr.ac.uk>

Brook Butler: brook.butler@srft.nhs.uk

Leicestershire, Northamptonshire and Rutland CLAHRC –

<http://clahrc-lnr.nihr.ac.uk>

Phil Shelton: pas35@leicester.ac.uk



IMPACT